

Animal welfare, etológia és tartástechnológia



Animal welfare, ethology and housing systems

Volume 9

Issue 3

Különszám/Special Issue

Gödöllő

2013

PATULIN – INDUCED CHANGES IN HAEMATOLOGICAL PARAMETERS OF RABBITS FED BY STRAWBERRY LEAVES AFTER CHRONIC EXPOSURE

Jana Emrichova^{1*}, Anna Kalafova¹, Katarina Zbynovska¹, Peter Petruska¹, Lubomir Ondruska², Rastislav Jurcik², Lubica Chrastinova², Anton Kovacik¹, Monika Schneidgenova¹, Peter Cupka¹, Marcela Capcarova¹

¹Department of Animal Physiology, Faculty of Biotechnology and Food Sciences, Slovak University of Agriculture in Nitra, Tr. A. Hlinku 2, 949 76 Nitra, Slovak Republic

²Animal Production Research Centre Nitra, Hlohovecka 2, 949 01 Nitra, Slovak Republic

* Corresponding author: e-mail: Jana Emrichova <emrichova@gmail.com>

INTRODUCTION

The flavonoids are a large group of naturally occurring compounds that are found in plants and are frequently consumed as part of the human diet. Flavonoids seem to play an important role in human health and to possess beneficial effects in the prevention of human diseases (Deepshikha *et al.*, 2008; Mareš *et al.*, 2008). Strawberry leaves have higher oxygen radical absorbance capacity (ORAC) than fruits (Shiow *et al.*, 2000). This leaves as phytogetic feed additive can improve welfare (Mareš *et al.*, 2008), restrict growth of fungi (Marcinčák *et al.*, 2010).

LITERATURE SURVEY

The strawberry leaves contain a wide range of phenolic compound (Hanhineva *et al.*, 2009), quercitrin, quercetin (Dreyer *et al.*, 2010; Oberbeil *et al.*, 2005), myricetin, kaempferol and epikatechin (Minárik, 2009).

Phytogetic feed additives are defined as a herbal substances included in the feed mixture for the purpose of enhancing production performance, improving performance feed and increase of quality animal products (Václavková *et al.*, 2010). Natural substances contained in these phytogetic feed additives affect (can improve) the sensory quality of feed (Mareš *et al.*, 2008). The use of plants in feed production can be restrict growth of fungi and production of mycotoxins in animals feed. They reduce the need of use drugs. Their feeding does not require of adherence to any withdrawal periods (Marcinčák *et al.*, 2010).

Mycotoxins are secondary metabolites of fungal origin (Painter *et al.*, 2003; Šimůnek, 2004; Serra *et al.*, 2005; Frisvad *et al.*, 2006). Patulin is produced by various species of *Aspergillus* and *Penicillium* (Polster, 1984; Betina, 1990; Malíř *et al.*, 2003; Frisvad *et al.*, 2007; González *et al.*, 2007). Patulin has bacteriostatic, bactericidal and fungicidal effects. It is toxic to plants and animals cells (Toman *et al.*, 2003; Sabater-Vilar, 2004), exhibits carcinogenic (Herzig, 2002; Sabater-Vilar, 2004), mutagenic and teratogenic activity (Sugiyanto *et al.*, 1993; Schumacher *et al.*, 2005), activate gastrointestinal disorders, anaemia, swelling and haemorrhage of various organs (Jesenská, 1987; Rimárová, 2002; Sabater-Vilar, 2004).

The aim of present study was to determinate the effect strawberry leaves inclusion to the feed mixture and single dose of patulin on haematological parameters of rabbits.

MATERIAL AND METHODS

Animals and diet

Fifteen adult male rabbits of Californian broiler line were used in experiment. Rabbits were obtained from an experimental farm of the Animal Production Research Centre in Nitra, Slovak Republic. Rabbits (in the age of 4 months, weighing 3.5 – 4.0 kg) were housed in individual flat-deck wire cages (area 0.34 m²). The animals were healthy and their condition was judged as good at the commencement of the experiment. Animals were kept in cages, at standard conditions (temperature

20 – 22°C, 14 h light period). Drinking water and feeding mixture for all animals was provided on an *ad libitum* basis. Animals were divided into four groups, one control group C (n =3) and three experimental groups E1, E2 and E3 (n = 4 in each group). Rabbits were fed with a granular feed mixture (FM) with strawberry leaves in various doses and all groups received patulin in injectable form at 10 µg.kg⁻¹ for 28 days 2 times a week (Table1).

Table 1.

Concentration of strawberry leaves and patulin of the experimental diet

Group	Concentration of strawberry leaves	Concentration of patulin (µg.kg ⁻¹)
Control (n = 3)	0 %	10
E1 (n = 4)	0.5 %	10
E2 (n = 4)	1.0 %	10
E3 (n = 4)	1.5 %	10

Blood sampling and analyses

Blood samples from *vena auricularis* were taken from all animals. In whole blood, selected haematological parameters as total white blood cell count (WBC), lymphocytes count (LYM), medium size cell count (MID), granulocytes count (GRA), lymphocyte percentage (LYM%), medium size cell percentage (MI%), granulocytes percentage (GRA%), red blood cell count (RBC), haemoglobin (HGB), haematocrit (HCT), mean corpuscular volume (MCV), mean corpuscular haemoglobin (MCH), mean corpuscular haemoglobin concentration (MCHC), red cell distribution width (RDWc), platelet count (PLT), platelet percentage (PCT), mean platelet volume (MPV) and platelet distribution width (PDWc) were measured using haematology analyzer Abacus junior VET (Diatron®, Vienna, Austria).

Statistical analyses

To compare the results, one-way ANOVA test was applied to calculate basic statistic characteristics and to determine significant differences between experimental and control groups. Statistical software SIGMA PLOT 11.0 (Jandel, Corte Madera, CA, USA) was used. Differences were compared for statistical significance at the level $P < 0.05$.

RESULTS AND DISCUSSION

The results of blood haematological parameters are summarized in Table 2. Addition of strawberry leaves and patulin influenced some haematological parameters in blood of rabbits. Statistical evaluation showed significant decrease of MCHC ($P < 0.05$) in E2 and E3 group in comparison with the experimental group E1. We observed significant lower values in HGB in E3 group in comparison with the control group. In contrast with results observed by *Petruška et al.* (2012), the values of HGB and MCHC were not influenced ($P > 0.05$) after quercetin/T-2 toxin treatment. Treatment with phytoadditives had no significant effect on HGB of male laboratory mice (*Singh et al.*, 2008) and rats (*Babayi et al.*, 2007; *Chinnadurai et al.*, 2013).

Decrease of HGB and MCHC in our study can be related with absence of iron and starting of anaemia (*Rolinec et al.*, 2010; *Bačovský*, 2013). Patulin caused the anaemia in study of *Camguilhem et al.* (1976) in sheep. The results of our study showed no significant differences in LYM, MID, RBC, HGB, HCT, MCH, PLT, PCT and MPV after long term application of patulin in combination with strawberry leaves. Selected haematological parameters were not influenced by quercetin and T-2 toxin in another study of *Petruška et al.* (2012).

Table 2.

Haematological parameters of rabbits after strawberry leaves and patulin treatment

Parameter	C	E1	E2	E3
WBC	12.97±1.21	13.53±2.58	13.61±2.25	10.39±2.28
LYM	8.22±2.52	7.39±4.72	7.48±1.98	6.82±1.3
MID	0.58±0.15	0.44±0.27	0.50±0.18	0.54±0.29
GRA	4.17±1.92	5.71±2.45	5.63±0.80	3.03±1.79
LY%	63.03±16.02	51.93±22.37	54.25±6.83	66.27±9.64
MI%	4.53±1.42	3.48±2.32	3.65±1.06	5.33±3.14
GR%	32.47±14.74	44.58±21.40	42.05±7.70	28.43±12.79
RBC	6.39±0.39	6.15±0.58	6.15±0.28	5.91±0.41
HGB	154.94±5.75 ^a	148.22±4.78	146.07±6.21	138.17±8.76 ^b
HCT	34.29±1.91	32.48±1.34	33.16±1.57	31.59±2.19
MCV	53.72±0.30	53.97±2.74	53.88±0.90	53.46±0.83
MCH	24.27±0.60	24.2±1.42	23.73±0.51	23.37±0.47
MCHC	452.13±8.80	456.46±4.35 ^a	440.7±12.67 ^b	437.53±3.54 ^b
RDWc	18.47±0.57	18.60±0.67	19.2±0.88	18.93±1.10
PLT	256.6±67.09	209.9±68.45	185.9±138.31	183.49±40.79
PCT	0.19±0.05	0.13±0.05	0.12±0.08	0.12±0.03
MPV	7.37±1.15	6.08±0.39	6.98±1.24	6.77±0.70
PDWc	33.27±0.61	30.08±2.10	33.85±4.87	32.57±3.27

^{a,b} – means in the same line with the different letters are different at the level $P < 0.05$

WBC - total white blood cell count ($10^9/l$); LYM - lymphocytes count ($10^9/l$); MID - medium-size cell count; GRA - granulocytes count ($10^9/l$); LYM% - lymphocyte percentage; MID% - medium-size cell percentage; GRA% - granulocytes percentage; RBC - red blood cell count ($10^{12}/l$); HGB - haemoglobin (g/l); HCT - haematocrit (%); MCV - mean corpuscular volume (fl); MCH - mean corpuscular haemoglobin (pg); MCHC - mean corpuscular haemoglobin concentration (g/l); RDWc - red cell distribution width (%); PLT - platelet count ($10^9/l$); PCT - platelet percentage; MPV - mean platelet volume (fl); PDWc - platelet distribution width (%), C – control group, E1, E2, E3 – experimental groups. The values shown are the mean ± SD (standard deviation).

CONCLUSIONS

Significant decrease of HGB and MCHC was probably caused by long term exposure of patulin, which can lead to anaemia. Selected concentrations of strawberry leaves in individual doses did not cause protection of the homeostasis. Higher concentrations could act as an antioxidant as we expected. Further experimental studies with strawberry leaves are needed to define the specific mechanisms of action.

Keywords: haematological parameters, strawberry leaves, patulin, rabbit, chronic exposure

Acknowledgments: This work was financially supported by VEGA scientific grant 1/0084/12, 1/0790/11, and KEGA grant 030SPU-4/2012.

REFERENCES

- Babayi, H. M., Udeme, J. J., Abalaka, J. A., Okogun, J. I., Salawu, O. A., Akumka, D. D., Adamu Zarma, S. S., Adzu, B. B., Abdulmumuni, S. S., Ibrahime, K., Elisha, B. B., Zakariys, S. S., Inyang, U. S. 2007. Effect of oral administration of aqueous whole extract of *Cassia filiformis* on haematograms and plasma biochemical parameters in rats. In *Journal of Medical Toxicology*. 2007, vol. 3, p. 146–151.
- Báčovský, J. (2013): Blood count interpretation. [online 5.8. 2013].
http://public.fnol.cz/www/3ik/vyuka/education_english/bacovsky/blood_count_interpretation_1.pdf
- Camguilhem, R., Escoula, L., Henry, M. (1976): Toxins of *Byssochlamys nivea* Westling. I. Preliminary study of toxicity in sheep. *Ann Rech Vet.* 1976;7(2):177-83.

- Chinnadurai, K., Kanwal, H.K., Tyagi, A.K., Stanton, C., Ross, P. (2013): High conjugated linoleic acid enriched ghee (clarified butter) increases the antioxidant and antiatherogenic potency in female Wistar rats. *Lipids Health Dis.* 2013 Aug 7;12(1):121.
- Deepshikha, M., Flora, S. J. S. (2008): Quercetin administration during chelation therapy protects arsenic-induced oxidative stress in mice. In *Biological Trace Element Research*. 2008, vol. 122, p. 137-147.
- Dreyer, E.M., Dreyer, W. (2010): Bylinky, plody a houby. Víkend, Český Tešín.
- Frisvad, J.C., Thrane, U. (2006): *Mycotoxin production by common filamentous fungi* [online 10.10. 2012]. <<http://www.cbs.knaw.nl/food/index.htm>>
- González, J., Osnaya, L., Soriano, J.M., Moltó, J.C., Mafles, J. (2007): Exposure assessment to patulin from the consumption of Apple-based products. In *Food additives and Contaminants*. no.24, p.11.
- Hanhineva, K., Soininen, P., Anttonen, J.M., Kokko, H., Rogachev, I., Aharoni, A., Laatikainen, R., Kärenlampia, S. (2009): NMR and UPLC-qTOF-MS/MS Characterisation of Novel Phenylethanol Derivatives of Phenylpropanoid Glucosides from the Leaves of Strawberry (*Fragaria A. ananassa* cv. Jonsok). In *Phytochem. Anal.*, vol. 20, p.353–364.
- Herzig, I. (2002): Mykotoxiny, jejich výskyt a vliv na zdraví. In *Krmivářství*. Výzkumný ústav veterinárního lékařství v Brně. no. 3, 2002, p. 11-12.
- Jesenská, Z. (1987): *Mikroskopické huby v požívatínách a v krmivách*. Bratislava : Alfa, p. 60-320. MHV 063-018-87.
- Malíř, F., Ostrý, V. (2003): *Vláknité mikromycety (plísňe), mykotoxiny a zdraví člověka*. Brno: Národní centrum ošetřovatelství a nelékařských zdravotnických oborů, p. 25-349. ISBN 80-7013-395-3.
- Marcinčák, S., Popelka, P., Martonová, M., Šimková, J. (2010): Vplyv rastlinných aditív na rastové parametre brojerových kurčiat. In *Krmivářství*, vol. 6, p. 11 – 12.
- Mareš, P., Zeman, L., Večerek, M. (2008): Využití fyto-genických přípravku ve výživě zvířat. In *Krmivářství*, vol. 1, p. 21 -23.
- Rolinec, M., Bíro, D., Šťastný, P., Kanka, T. (2010): Analyzis of haematological profile of piglets. In *Acta fytotechnica et zootechnica*. Nitra, Slovaca Universitas Agriculturae Nitriae, 2010, p. 40-43.
- Minárik, P. (2009): Flavonoidy. *Bedeker zdravia*, vol. 2, p.70 -71.
- Oberbeil, K., Lentzová, CH. (2005): *Ovocie a zelenina ako liek*. Fortuna Print. Bratislava.
- Petruška, P., Capcarová, M. (2012): Effect of chronic application of quercetin and acute dose of T-2 toxin on haematological parameters of rabbits : Vplyv chronickej aplikácie kvercetínu a akútnej dávky T-2 toxínu na hematologické parametre kráľika. In *MendelNet 2012*. Brno: Mendelova univerzita, 2012, p. 981--988. ISBN 978-80-7375-563-8.
- Rimárová, K. (2002): Karcinogénne látky v životnom prostredí a potravinovom reťazci. In *Životné prostredie*, vol. 36, no. 3, p. 129-132.
- Sabater-Vilar, M. (2004): Patulin produced by an *Aspergillus clavatus* isolated from feed containing malting residues associated with a lethal neurotoxicosis in cattle. In *Mycopathologia*, vol. 158, no. 4, p. 419-426.
- Schumacher, D.M., Metzler, M., Lehmann, L. (2005): Mutagenicity of the mycotoxin patulin in cultured Chinese hamster V79 cells, and its modulation by intracellular glutathione. In *Arch Toxicol*, vol. 79, p.110–121.
- Serra, R., Braga, A., Venancio, A. (2005): Mycotoxin-producing and other fungi isolated from grapes for wine production, with particular emphasis on ochratoxin A. In *Research in Microbiology*, vol. 156, 2005, no. 4, p. 515 – 521.
- Shiow Y. Wang, Hsin-Shan Lin. (2000): Antioxidant Activity in Fruits and Leaves of Blackberry, Raspberry, and Strawberry Varies with Cultivar and Developmental Stage. In *J. Agric. Food Chem.*, vol. 48, no. 2, p. 140–146.
- Šimůnek, J. (2004): *Plísňe a mykotoxiny*. Brno. 2004. [cit. 2012-10-10] http://www.med.muni.cz/dokumenty/pdf/plisne_a_mykotoxiny.pdf.
- Singh, A., Singh, S. K. (2008): Reversible antifertility effect of aqueous leaf extract of *Allamanda cathartica* L. in male laboratory mice. In *Andrologia*, vol. 40, 2008, p. 337–345.
- Sugiyanto, J., Inouye, M., Oda, S.I., Takagishi, Y., Yamamura, H. (1993): Teratogenicity of patulin, a mycotoxin, in mice. In *Environ. Med.* 37, p. 43–46.
- Toman, R., Golian, J., Massányi, P. (2003): *Toxikológia potravín*. Nitra : SPU v NITRE, p. 62-65. ISBN 80-8069-166-5.
- Václavková, E., Lustyková, A. (2010): Fyto-geninní krmná aditiva ve výživě monogastru. In *Krmivářství*, vol. 6, p. 9 – 10.