

Animal welfare, etológia és tartástechnológia



Animal welfare, ethology and housing systems

Volume 9

Issue 3

Különszám/Special Issue

Gödöllő

2013

ASSESSMENT OF BENEFICIAL AND TOXIC EFFECTS OF NATURAL SUBSTANCES ON OVARIAN FUNCTIONS

Adriana Kolesárová*¹, Nora Maruniaková¹, Marina Medved'ová¹, Marek Halenár¹, Attila Kádasi^{1,2}, Andreia Filipa Tomás Marques³, Jozef Bulla¹

¹Department of Animal Physiology, Faculty of Biotechnology and Food Sciences, Slovak University of Agriculture in Nitra, Nitra, Slovak Republic;

²Institute for Genetics and Reproduction of Farm Animals, Animal Production Research Centre Nitra, Lužianky, Slovak Republic;

³Tropical Research Institute, Lisboa, Portugal,

*Corresponding author: Adriana.Kolesarova@uniag.sk

ABSTRACT

The dynamics of a stress response of the organism towards potential stressors – mycotoxins underlines the need to define limits in which animals develop a state without compromising their health and maintain a stable internal milieu. Tissue and cell cultures are of increasing interest in the evaluation of toxicological risks of toxic compounds and their possible elimination. It is for this reason the solution of the problem in the field of examining effects of the potential protective natural substances in the animal organism and their mechanism of the effect is highly relevant. Ovarian granulosa cells are sensitive to the effects of mycotoxin – trichothecenes but bioactive substances of natural origin may be important in the prevention of reproductive alterations induced by exogenous and endogenous factors.

Keywords: ovary, deoxynivalenol, resveratrol, steroidogenesis, proliferation, apoptosis

INTRODUCTION

The environmental stress is one of the main factors that change the reproductive functions (Kolesarova et al., 2010a,b,c; 2011a,b; 2012). Previous reports describe the influence of beneficial (Kolesarova et al., 2011c, 2012) and toxic (Kolesarova et al., 2009; 2010a,b,c; 2011a,b; Medvedova et al., 2011) substances on porcine ovarian (GCs) focused on processes of steroidogenesis, proliferation and apoptosis in these reproductive cells (Kolesarova et al., 2009; 2010a,b,c; 2011a,b; Medvedova et al., 2011). On the other hand the bioactive substances of natural origin may be important in the prevention of reproductive alterations induced by exogenous and endogenous factors (Kolesarova et al., 2012).

Characteristic of Trichothecenes - Deoxynivalenol

Mycotoxins are natural and very stable toxins, with relatively low-molecular weight secondary metabolites of fungal origin, which can contaminate a large variety of feed mixtures (Labuda et al., 2009; Tancinova et al., 2009), grains and foodstuffs worldwide (Schollenberger et al., 2007; Ranzenigo et al., 2008). They are found in a variety of foods and beverages, including both plant-based products and animal products. Among the first ones, its presence in cereal grains (corn, wheat, barley, oats, rye, rice, etc.) and beans (coffee, cocoa, soy, etc.) are harmful to animals and humans (Abouzied et al., 1991). However, not all fungi produce mycotoxins and among the toxigenic species, some only produce one type of mycotoxin, while others are able to produce several. Cereals can be mostly contaminated by *Fusarium* species, a group of toxin-producing molds (Tiemann et al., 2003a,b; Larsen et al., 2004; Giraud et al., 2010).

Trichothecenes, such as deoxynivalenol (DON), are the major mycotoxins of *Fusarium species* (Larsen et al., 2004; D'Mello et al., 1999). DON (C₁₅H₂₀O₆; mol. wt. = 296.32) is also called Dehydronivalenol, 4-Deoxynivalenol, 4-Desoxynivalenol or Vomitoxin. DON can be acutely or

chronically toxic, or both, depending on the kind of toxin and the dose. Also, it could be rapidly absorbed after oral administration passively throughout the gastrointestinal tract and actively in the kidneys (Marquardt and Frohlich, 1992), liver, muscle, adipose tissue (Gareis et al., 2000) and reproductive tissues (Ranzenigo et al., 2008). Major *Fusarium* mycotoxin, DON, negatively influences the establishment of pregnancy in pigs in part because of its ability to inhibit oocyte maturation (Alm et al., 2002; 2006). Thus, mycotoxin exposure that alters granulosa cells and steroid hormone production may also alter oocyte development (Alm et al., 2002), ovulation (Kumagai et al., 1982), reproductive functions and pregnancy outcome (D'Mello et al., 1999; Alm et al., 2002).

Mechanism of the effect of Trichothecenes - Deoxynivalenol

Trichothecene toxicity at the cellular level is characterized by inhibited protein synthesis (Rotter et al., 1994), impairment of membrane functions, altered intercellular communication and deregulation of calcium homeostasis (Pestka and Smolinski 2005). Trichothecenes not only bind eukaryotic ribosomes and interfere with translation (Ueno 1983), but also activate intracellular protein kinases that both mediate selective gene expression and apoptosis, ultimately contributing to downstream pathologic sequelae (Pestka et al., 2004). Moreover, it has been demonstrated that trichothecenes rapidly activate mitogen-activated protein kinase (MAPK) which modulates physiological processes including cell growth, differentiation and apoptosis (Kouadio et al., 2007, Marzocco et al., 2009; Luongo et al., 2010). In addition to inhibiting translation, trichothecenes can simultaneously activate p38, Jun N-terminal Kinase (JNK) and extracellular signal-regulated kinase (ERK), MAPK *in vitro* and *in vivo* (Chung et al., 2003; Moon and Pestka, 2003; Moon et al., 2003; Yang et al., 2000; Zhou et al., 2003a,b, 2005a,b) via a process referred to as "ribotoxic stress" (Iordanov et al., 1997). Oxidative stress is certainly involved in the toxicities of trichothecene mycotoxins (El Golli et al., 2006). Oxidative stress in the form of reactive oxygen species (ROS) generation or disruption of the redox balance in the cell not only induces apoptosis but also involved in cell proliferation and signalling (Martindale and Holbrook, 2002). Oxidative stress is the underlying mechanism by which trichothecene causes DNA damage and apoptosis *v.* At the cellular level, DON induces ribotoxic stress thereby disrupting macromolecule synthesis, cell signaling, differentiation, proliferation, and death (Pestka, 2010).

Trichothecenes affect ovarian functions

Secretory activity: The influence of mycotoxin - DON on secretory activity was detected by Medvedova et al. (2011) and Kolesarova et al. (2012). Dose-response of DON on ovarian granulosa cells was examined in these studies. Isolated ovarian GCs were able to survive in culture and release hormonal substances insulin-like growth factor I (IGF-I) and progesterone after experimental DON addition at the doses 10, 100 and 1000 ng/ml. These data confirm previous reports concerning the influence of toxic substances on porcine cellular processes (Ranzenigo et al., 2008; Kolesarova et al., 2009, 2010a,b,c,d; 2011a,b). The effect of DON in relation to IGF-I release by ovarian granulosa cells have been examined previously by Ranzenigo et al. (2008) but the time of cell culture with mycotoxin application lasted 2 days and in the study of Medvedova et al. (2011) it was 24 h. DON had inhibitory effects on IGF-I –induced steroid production and decreased cell numbers at the dose of 1000 ng/ml (Ranzenigo et al., 2008). Medvedova et al. (2011) have shown that the mycotoxin DON decreased IGF-I secretion by cultured GCs at a dose of 1000 ng/ml but it was not affected by the doses of 10 ng/ml and 100 ng/ml. Ranzenigo et al. (2008) have demonstrated inhibited progesterone production induced by FSH plus IGF-I at the doses 100 ng/ml (0.337 μ M) and 1000 ng/ml (0.37 μ M) of DON. In the study of Medvedova et al. (2011) visible mitosis of GCs after 24 h DON effect at the dose of 10 ng/ml was observed. Progesterone release was stimulated by DON at the dose 1000 ng/ml but not at 10 and 100 ng/ml. (Medvedova et al., 2011). Similarly progesterone release was stimulated by toxic doses of DON (2000, 3000 and 5000 ng/ml) used in the experiment (Kolesarova et al., 2012). The effect of the DON was found to be concentration dependent as it was described in previous study (Ranzenigo et al., 2008; Medvedova et al., 2011; Kolesarova et al., 2012).

Proliferation and apoptosis: Expressions of proliferation (cyclin B1 and PCNA)- associated peptides were stimulated by DON (Medvedova et al., 2011). Analyses of porcine endometrial cells cultivated (for 24h) with different doses of DON were done by Tiemann (2003a). These results point out to significant decrease of proliferation at the concentrations of 1.88 μM and 3.76 μM of DON. These authors also observed expression of PCNA protein, which significantly decreased at the doses of 1.88 μM (for 24h) and 0.94 μM (for 48h) of DON. In the study of Medvedova et al. (2011) expression of cyclin B1 was stimulated by DON (at 1000 ng/ml but not at 10 and 100 ng/ml). Similarly PCNA expression was stimulated by DON (at 100 and 1000 ng/ml but not at 10 ng/ml). On the other hand caspase-3 expression was not influenced by DON treatment (at doses 10, 100 and 1000 ng/ml) (Medvedova et al., 2011).

Conclusion: Porcine ovarian GCs are sensitive to the effects of mycotoxin - trichothecenes. The results indicate, (1) a direct effect of DON on secretion of hormones, (2) expression of markers of proliferation (cyclin B1 and PCNA) but not (3) expression of marker of apoptosis (caspase-3) in porcine ovarian granulosa cells. This in vitro study suggests the dose-dependent association of DON on porcine ovarian functions (Medvedova et al., 2011).

Characteristic of Phytoalexins- Resveratrol

Phytoalexins are antimicrobial substances under natural conditions not found in plants. Their creation occurs in plants attacked by pathogens (Dixon and Paiva, 1995). Resveratrol (RSV, 3,5,4'-trihydroxystilbene) was the first isolated from the roots of white hellebore (*Veratrum grandiflorum* O. Loes) in 1940 (Takaoka, 2006) and later in 1963, from the roots of (*Polygonum cuspidatum*) a plant used in traditional Chinese and Japanese medicine (Nonomura et al., 1963). It is a natural polyphenol widely present in plants and in particular in the skin of red grapes and in wine, resveratrol antioxidant properties have been well demonstrated, with a wide range of biological effects (Karuppagounder et al., 2009). It is beneficial against diverse cardiac diseases including ischemic heart disease, hypertrophy, heart failure, atherosclerosis, hypertension, diabetes and obesity (Bertelli and Das, 2009; Mukherjee et al., 2009). Resveratrol is well known for its phytoestrogenic, antioxidant properties, growth-inhibitory and apoptosis-inducing activities (Joe et al., 2002; Jiang et al., 2005; Baur and Sinclair, 2006). It also has protective role in endothelial cells by modulating mitochondrial oxidative stress (Ungvari et al., 2009).

Mechanism of the effect of Phytoalexins- Resveratrol

Resveratrol is able to inhibit PI3K/Akt Protein Kinase/ mammalian target of rapamycin (PI3K/Akt/mTOR) pathway in various types of cells (Jiang et al., 2005). It inhibits proliferation and induces apoptosis in cells at different levels (Ferry-Dumazet et al, 2002, Haider et al, 2003). Resveratrol can alter a variety of genes thereby changing the "death signal" into a "survival signal" (Das and Maulik, 2004). Svechnikov et al. (2009) present investigation has demonstrated that resveratrol and its analogues structure-dependently attenuated steroidogenesis in Leydig cells through suppression of the expression of StAR and cytochrome P450c17.

Phytoalexins affect ovarian functions

Secretory activity: Stimulatory effect of RSV on progesterone release by GCs was recorded after resveratrol treatment at the dose of 50 $\mu\text{g/ml}$, while doses of 30 and 10 $\mu\text{g/ml}$ did not affect the release of the steroid hormone (Kolesarova et al., 2012). RSV in combination with DON at the highest doses (50 $\mu\text{g/ml}$ of RSV and 5000 ng/ml of DON) stimulated progesterone release by GCs. Stimulatory effect of alone DON on the progesterone release by GCs was lower in comparison with RSV in combination with DON (Kolesarova et al., 2012). The effect of the DON (Medvedova et al., 2011; Ranzenigo et al., 2008) and DON combined with resveratrol (Kolesarova et al., 2012) was found to be concentration dependent.

Conclusion: The results of Kolesarova et al. (2012a) indicate, (1) the dose-dependent stimulatory effects of RSV, DON and combination of RSV with DON on release of steroid hormone progesterone and (2) reduction of the stimulatory effect of DON by RSV. The *in vitro* results suggest that reproductive toxicity of animals induced by mycotoxin - deoxynivalenol can be inhibited by protective natural substance – resveratrol.

Conclusion

In conclusion, it is important to highlight that the impact of fungal toxins upon animals extends beyond their clinical features. Despite the ever-increasing understanding of mycotoxins, they still have a continuous and severe economic impact worldwide (Marquardt, 1996; Hussein and Brasel, 2001). Although acute ingestion of high levels of mycotoxins can be very harmful to the animal, long term consumption of low concentrations of mycotoxins can also be damaging. There are several ways of reducing mycotoxin concentrations both pre and post-harvest, including the addition of feed additives such as, but not limited to, natural clays, yeasts, and enzymes. Since mycotoxins can be so detrimental to swine industry, further determination of sustainable way to combat the global mycotoxin problem is important for maintaining animal health, as well as reducing economic impacts on farmers and producers (Chaytor et al., 2011). Tissue and cell cultures are of increasing interest in the evaluation of toxicological risks of contaminated compounds and their possible elimination. It is for this reason the solution of the problem in the field of examining effects of the potential protective natural substances in the animal organism and their mechanism of the effect is highly relevant.

Acknowledgments:

The results shown in this review were obtained financially supported by the Ministry of Education, Science, Research and Sport of the Slovak Republic projects no. 1/0790/11, 1/0022/13 and APVV-0304-12, KEGA 030SPU-4/2012, RGB Net (COST Action TD1101), European Community under project no 26220220180: Building Research Centre „AgroBioTech”.

REFERENCES

- Abouzied, M.M., Azcona, J.I., Braselton, W.E., Pestka, J.J. (1991): Immunochemical assessment of mycotoxins in 1989 grain foods: Evidence for deoxynivalenol (vomitoxin) contamination. *Appl. Environ. Microbiol.* 1991, 57, 3672–3677.
- Alm, H., Greising, T., Brüßow, K.P., Torner, H., Tiemann, U. (2002): The influence of the mycotoxins deoxynivalenol and zearalenol on *in vitro* maturation of pig oocytes and *in vitro* culture of pig zygotes. *Toxicol. In Vitro*, 2002, 16, 643–648.
- Alm, H., Brüßow, K.P., Torner, H., Vanselow, J., Tomek, W., Danicke, S., Tiemann, U. (2006): Influence of Fusarium-toxin contaminated feed on initial quality and meiotic competence of gilt oocytes. *Reprod. Toxicol.*, 2006, 22, 44–50.
- Baur, J.A., Sinclair, D.A. (2006): Therapeutic potential of resveratrol: the *in vivo* evidence. *Nature reviews Drug Discovery*, 5, 493-506.
- Bertelli, A.A., Das, D.K. (2009): Grapes, wines, resveratrol, and heart health. *J. Cardiovasc. Pharmacol.* 2009, 54, 468–476.
- Chaudhari, M., Jayaraj, R., Bhaskar, A.S., Lakshmana Rao, P.V. (2009): Oxidative stress induction by T-2 toxin causes DNA damage and triggers apoptosis via caspase pathway in human cervical cancer cells. *Toxicology*, 2009, 262(2), 153-61.
- Chaytor, A.C., Hansen, J.A., Heugten, E., Todd See, M., Woo Kim, S. (2011): Occurrence and decontamination of mycotoxins in swine feed. *Asian-Aust. J. Anim. Sci.*, 2011, 24(5), 723-738.
- Chung, Y.J., Zhou, H.R., Pestka, J. (2003): Transcriptional and post-transcriptional roles for p38 mitogen-activated protein kinase in upregulation of TNF- α expression by deoxynivalenol (vomitoxin). *Toxicol. Appl. Pharmacol.*, 2003, 193, 188–201.
- Das, D.K., Maulik, N. (2004): Conversion of death signal into survival signal by redox signaling. *Biochemistry (Mosc)*, 2004, 69, 10–17.
- Dixon, R.A., Paiva, N.L. (1995): Stress-induced phenylpropanoid metabolism. *Plant Cell*, 7, 1085-1097.

- D'Mello, J.P.F., Placinta, C.M., Macdonald, A.M.C. (1999): Fusarium mycotoxins, a review of global implications for animal health, welfare and productivity, *Anim. Feed Sci. Technol.*, 1999, 80, 183–205.
- El Golli, E., Hassen, W., Bouslimi, A., Bouaziz, C., Ladjimi, M.M., Bacha, H. (2006): Induction of Hsp 70 in Vero Cells in Response to Mycotoxins Cytoprotection by Sub-Lethal Heat Shock and by Vitamin E. *Toxicol. Lett.*, 2006, 166, 122-130.
- Ferry-Dumazet, H., Garnier, O., Mamani-Matsuda, M., Vercauteren, J., Belloc, F., Billiard, C., Dupouy, M., Thiolat, D., Kolb, J. P., Marit, G., Reiffers, J., Mossalayi, M. D. (2002): Resveratrol inhibits the growth and induces the apoptosis of both normal and leukemic hematopoietic cells. *Carcinogenesis*, 23, 8, 1327-1333.
- Gareis, M., Wolff, J. Relevance of mycotoxin contaminated feed for farm animals and carryover of mycotoxins to food of animal origin. *Mycoses*, 2000, 43 Suppl 1:79-83.
- Giraud, F., Pasquali, M., El Jarroudi, M., Vrancken, C., Brochot, C., Cocco, E., Hoffmann, L., Delfosse, P., Bohn, T. (2010): Fusarium head blight and associated mycotoxin occurrence on winter wheat in Luxembourg in 2007/2008. *Food Addit. Contam. Part A, Chem. Anal. Control Expo. Risk. Assess.*, 2010, 26, 1–11.
- Haider, U. G., Sorescu, D., Griendling, K. K., Vollmar, A. M., Dirsch, V. M. (2003): Resveratrol increases serine15-phosphorylated but transcriptionally impaired p53 and induces a reversible DNA replication block in serum-activated vascular smooth muscle cells. *Molecular Pharmacology*, 2003, 63, 4, 925-932.
- Hussein, H.S., Brasel, J.M., 2001: Toxicity, metabolism, and impact of mycotoxins on humans and animals. *Toxicology*, 2001, 167, 101–134.
- Iordanov, M.S., Pribnow, D., Magun, J.L., Dinh, T.H., Pearson, J.A., Chen, S.L., Magun, B.E. (1997): Ribotoxic stress response: activation of the stress-activated protein kinase JNK1 by inhibitors of the peptidyl transferase reaction and by sequence-specific RNA damage to the alpha-sarcin/ricin loop in the 28S rRNA. *Mol Cell Biol.*, 1997, 17(6), 3373–3381.
- Jiang, H., Zhang, L., Kuo, J., Kuo, K., Gautam, S.C., Groc, L., Rodriguez, A.I, Koubi, D., Hunter, T.H., Corcoran, G.B., Seidman, M.D. Levine, R.L. (2005). Resveratrol-induced apoptotic death in human U251 glioma cells. *Molecular Cancer Therapy*, 4, 554-561.
- Joe, A.K, Liu, H., Suzui, M., Vural, M.E., Xiao, D., Weinstein, I.B. (2002): Resveratrol induces growth inhibition, S-phase arrest, apoptosis, and changes in biomarker expression in several human cancer cell lines. *Clinical Cancer Research*, 8, 893-903.
- Karuppagounder, S.S., Pinto, J.T., Xu, H., Chen, H.L., Beal, M.F., Gibson, G.E. (2009): Dietary supplementation with resveratrol reduces plaque pathology in a transgenic model of Alzheimer's disease. *Neurochem Int.*, 2009, 54, 111–118.
- Kolesarova, A., Slivkova, J., Sirotkin, A., Massanyi, P., Capcarova, M. (2009): The release of insulin-like growth factor-I by ovarian granulosa cells of pregnant sows after lead and mercury administration in vitro. *Slovak J. Anim. Sci.*, 2009, 42, 35–41.
- Kolesarova, A., Roychoudhury, S., Slivkova, J., Sirotkin, A., Capcarova, M., Massanyi, P. (2010a): In vitro study on the effect of lead and mercury on porcine ovarian granulosa cells. *Journal of Environmental Science and Health, Part A Toxic hazardous substances and environmental engineering*, 2010a, 45, 320-331.
- Kolesarova, A., Capcarova, M., Sirotkin, A., Medvedova, M., Kovacic, J. (2010b): Cobalt-induced changes in the IGF-I and progesterone release, expression of proliferation- and apoptosis-related peptides in porcine ovarian granulosa cells in vitro. *Journal of Environmental Science and Health, Part A*, 2010b, 45, 810-817.
- Kolesarova, A., Sirotkin, A., Roychoudhury, S., Capcarova, M. (2010c): Puberty related changes in hormonal levels, productive performance, carcass traits, and their interactions in Slovakian White gilts. *Asian–Australian Journal of Animal Science*, 2010c, 23, 182-187.
- Kolesarova, A., Capcarova, M., Roychoudhury, S. (2010d): Metal induced ovarian signaling. - Nitra : Slovak Agricultural University, 2010d, 135p. ISBN 978-80-552-0456-7.
- Kolesarova, A., Capcarova, M., Medvedova, M., Sirotkin, A.V., Kovacic, J. (2011a) In vitro assessment of iron effect on porcine ovarian granulosa cells: secretory activity, markers of proliferation and apoptosis. *Physiol Res.*, 2011a, 60(3), 503-10.
- Kolesarova, A., Capcarova, M., Sirotkin, A.V., Medvedova, M., Kovacic, J. (2011b): In vitro assessment of silver effect on porcine ovarian granulosa cells. *J Trace Elem Med Biol.*, 2011b, 25(3), 166-70.
- Kolesarova, A., Capcarova, M., Bakova, Z., Galik, B., Juracek, M., Simko, M., Sirotkin, A.V. (2011c) The effect of bee pollen on secretion activity, markers of proliferation and apoptosis of porcine ovarian granulosa cells in vitro. *J Environ Sci Health B*. 2011c, 46(3), 207-12.
- Kolesarova, A.; Capcarova, M.; Maruniakova, N.; Lukac, N.; Ciereszko, R. E.; Sirotkin, A.V. (2012): Resveratrol inhibits reproductive toxicity induced by deoxynivalenol. In *J Environ Science and Health, part A*, 2012, 47(9), 329-34.

- Kouadio, J. H., Sebastien, D. D., Moukha, S. (2007): Effect of combinations of fusarium mycotoxins on the inhibition of macromolecular synthesis malondialdehyde levels, DNA methylation and fragmentation and viability in Caco-2 cells. *Toxicon*, 2007, 49, 306–317.
- Kumagai, S., Shimizu, T. (1982): Neonatal exposure to zearalenone causes persistent anovulatory estrus in the rat. *Arch. Toxicol.*, 1982, 50, 279–286.
- Labuda, R., Parich, A., Vekiru, E., Tancinova, D. (2009): Incidence of fumonisins, moniliformin and Fusarium species in poultry feed mixtures from Slovakia. *Ann. Agric. Environ. Med.*, 2009, 12(1), 81–86.
- Larsen, J.C., Hunt, J., Perrin, I., Ruckebauer, P. (2004): Workshop on trichothecenes with a focus on DON: summary report. *Toxicol. Lett.*, 2004, 153(1), 1–22.
- Luongo, D., Severino, L., Bergamo, P., D'ariento, R., Rossi, M. (2010): Trichothecenes NIV and DON modulate the maturation of murine dendritic cells. *Toxicon.*, 2010, 55, 73–80.
- Marquardt, R.R., Frohlich, A.A. (1992): A review of recent advances in understanding ochratoxicosis. *Journal of Animal Science*, 1992, 70, 3968–3988.
- Marquardt, R.R. 1996. Effects of molds and their toxins on livestock performance: a western Canadian perspective. *Animal Feed Sci. Technol.*, 1996, 58, 77–89.
- Martindale, J.L., Holbrook, N.J. (2002): Cellular response to oxidative stress: signaling for suicide and survival. *J. Cell Physiol.*, 2002, 192, 1–15.
- Marzocco, S., Russo, R., Bianco, G., Autore, G., Severino, L. (2009): Pro-apoptotic effects of nivalenol and deoxynivalenol trichothecenes in J774A.1 murine macrophages. *Toxicol. Lett.*, 2009, 189, 21–26.
- Medvedova, M., Kolesarova, A., Capcarova, M., Labuda, R., Sirotkin, A.V., Kovacik, J., Bulla, J. (2011): The effect of deoxynivalenol on the secretion activity, proliferation and apoptosis of porcine ovarian granulosa cells in vitro. *J. Environ. Sci. Health B*, 2011, 46(3), 213–219.
- Moon, Y., And Pestka, J.J. (2003a): Deoxynivalenol-induced mitogen-activated protein kinase phosphorylation and IL-6 expression in mice suppressed by fish oil. *J. Nutr. Biochem.*, 2003a, 14, 717–726.
- Moon, Y., Uzarski, R., Pestka, J. (2003b): Relationship of trichothecene structure to COX-2 induction in the macrophage: selective action of Type B (8-keto) trichothecenes. *J. Toxicol. Environ. Health*, 2003b, 66, 1967–1983.
- Mukherjee, S., Lekli, I., Gurusamy, N., Bertelli, A.A., Das, D.K. (2009): Expression of the longevity proteins by both red and white wines and their cardioprotective components, resveratrol, tyrosol, and hydroxytyrosol. *Free Radic. Biol. Med.*, 2009, 46, 573–578.
- Nonomura, S., Kanagawa, H., Makimoto, A. (1963): Chemical constituents of polygonaceous plants. I. Studies on the components of Ko-jo-kon (*Polygonum cuspidatum* Sieb. Et Zucc.). *Yakugaku Zasshi*, 1963, 83, 988–990.
- Pestka, J., Zhou, H.R., Moon, Y., Chung, Y.J. (2004): Cellular and molecular mechanisms for immune modulation by deoxynivalenol and other trichothecenes: unraveling a paradox. *Toxicology Letters*, 2004, 153, 61–73.
- Pestka, J.J., Smolinski, A.T. (2005): Deoxynivalenol: toxicology and potential effects on humans. *J. Toxicol. Environ. Health B Crit. Rev.*, 2005, 8(1), 39–69.
- Pestka, J.J. (2010): Deoxynivalenol: mechanisms of action, human exposure, and toxicological relevance. *Arch. Toxicol.*, 2010, 84(9), 663–679.
- Ranzenigo, G., Caloni, F., Cremonesi, F., Aad, P.Y., Spicer, L.J. (2008): Effects of Fusarium mycotoxins on steroid production by porcine granulosa cells. *Anim. Reprod. Sci.*, 2008, 107(1–2), 115–130.
- Rotter, B.A., Thompson, B.K., Lessard, M., Trenholm, H.L., Tryphonas, H. (1994): Influence of low-level exposure to Fusarium mycotoxins on selected immunological and hematological parameters in young swine. *Fundam. Appl. Toxicol.*, 1994, 23, 117–124.
- Schollenberger, M., Muller, H.M., Ruffe, M., Terry-Jara, H., Suchy, S., Plank, S., Drochner, W. (2007): Natural occurrence of Fusarium toxins in soy food marketed in Germany. *Int. J. Food Microbiol.*, 2007, 113, 142–146.
- Svechnikov, K., Spatafora, C., Svechnikova, I., Tringali, C., Soder, O. (2009): Effects of resveratrol analogs on steroidogenesis and mitochondrial function in rat Leydig cells in vitro. *J. Appl. Toxicol.*, 2009, 29, 673–680.
- Takaoka, M.J. (1940): Of the phenolic substances of white hellebore (*Veratrum grandiflorum* Loes. Fil.). *J. Faculty Sci. Hokkaido Imperial University*, 1940, 3, 1–16.
- Tancinova, D., Labuda, R. (2009): Fungi on wheat bran and their toxinogenicity. *Ann. Agric. Environ. Med.*, 2009, 16(2), 325–331.
- Tiemann, U., Tomek, W., Schneider, F., Vanselow, J. (2003a): Effects of the mycotoxins alpha- and beta-zearalenol on regulation of progesterone synthesis in cultured granulosa cells from porcine ovaries. *Reprod. Toxicol.*, 2003a, 17(16), 673–681.
- Tiemann, U., Viergutz, T., Jonas, L., Schneider, F. (2003b): Influence of the mycotoxins α - and β -zearalenol and deoxynivalenol on the cell cycle of cultured porcine endometrial cells. *Reproductive Toxicology*, 2003b, 17, 209–218.

- UENO, Y. (1983): General toxicology. In: Ueno, Y. (Ed.), *Developments in food science. Trichothecenes. Chemical, biological and toxicological aspects.* Elsevier, Tokyo, 135–146.
- Ungvari, Z., Labinsky, N., Mukhopadhyay, P., Pinto, J.T., Bagi, Z., Ballabh, P., Zhang, C., Pacher, P., Csiszar, A. (2009): Resveratrol attenuates mitochondrial oxidative stress in coronary arterial endothelial cells. *Am. J. Physiol. Heart Circ. Physiol.*, 2009, 297, 1876–1881.
- Yang, G., Jarvis, B.B., Chung, Y., Pestka, J.J. (2000): Apoptosis induction by the satratoxins and other trichothecene mycotoxins: relationship to ERK, p38 MAPK and SAPK/JNK activation. In *Toxicol. Appl. Pharmacol.*, 2000, 164, 149–160.
- Zhou, H.R., Islam, Z., Pestka, J.J. (2003a): Rapid, sequential activation of mitogen-activated protein kinases and transcription factors precedes proinflammatory cytokine mRNA expression in spleens of mice exposed to the trichothecene vomitoxin. *Toxicol. Sci.*, 2003a, 72, 130–142.
- Zhou, H.R., Lau, A.S., Pestka, J.J. (2003b): Role of double-stranded RNA-activated protein kinase (PKR) in deoxynivalenol-induced ribotoxic stress response. *Toxicol. Sci.*, 2003b, 74, 335–344.
- Zhou, H.R., Islam, Z., Pestka, J.J. (2005a): Induction of competing apoptotic and survival signaling pathways in the macrophage by the ribotoxic trichothecene deoxynivalenol. *Toxicol. Sci.*, 2005a, 87(1), 113–122.
- Zhou, H.R., Jia, Q., Pestka, J.J. (2005b): Ribotoxic stress response to the trichothecene deoxynivalenol in the macrophage involves the SRC family kinase Hck. *Toxicol. Sci.*, 2005b, 85 (2), 916–926.